

Muscle metaboreflex and cerebral blood flow regulation in humans

Prodel, Eliza; Balanos, George M; Braz, Igor D; Nóbrega, Antonio Claudio L; Vianna, Lauro C; Fisher, James P

DOI:

[10.1152/ajpheart.00894.2015](https://doi.org/10.1152/ajpheart.00894.2015)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Prodel, E, Balanos, GM, Braz, ID, Nóbrega, ACL, Vianna, LC & Fisher, JP 2016, 'Muscle metaboreflex and cerebral blood flow regulation in humans: implications for exercise with blood flow restriction', *American Journal of Physiology Heart and Circulatory Physiology*. <https://doi.org/10.1152/ajpheart.00894.2015>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked March 2016

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Muscle metaboreflex and cerebral blood flow regulation in humans: implications for exercise with blood flow restriction

Eliza Prodel^{1,2}, George M. Balanos¹, Igor D. Braz¹, Antonio C. L. Nobrega², Lauro C. Vianna³ & James P. Fisher¹

¹School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences, University of Birmingham, Birmingham, UK; ²Department of Physiology and Pharmacology, Fluminense Federal University, Niterói, Brazil; ³Faculty of Physical Education, University of Brasília, DF, Brazil

Running Title: Cerebral blood flow in dynamic exercise

Please send correspondence to:

James P. Fisher, Ph.D.

School of Sport, Exercise and Rehabilitation Sciences

College of Life and Environmental Sciences

University of Birmingham

Edgbaston, Birmingham

B15 2TT, England

Tell: +44 (0)121 414 8011

Fax: +44 (0)121 414 4121

Email: j.p.fisher@bham.ac.uk

ABSTRACT

We investigated the effect of activating metabolically sensitive skeletal muscle afferents (muscle metaboreflex) on cerebral blood flow and the potentially confounding influence of concomitant changes in the partial pressure of arterial carbon dioxide. Eleven healthy males (25 ± 4 years) performed submaximal leg cycling exercise on a semi-recumbent cycle ergometer (heart rate ~ 120 b \cdot min $^{-1}$), and assessments made of the partial pressure of end-tidal carbon dioxide ($P_{ET}CO_2$), internal carotid artery blood flow (ICA_Q) and conductance (ICA_{CVC}), middle cerebral artery mean blood velocity (MCA_{Vm}) and conductance index (MCA_{CVCi}). The muscle metaboreflex was activated during cycling with leg blood flow restriction (BFR) or isolated with post exercise ischemia (PEI). In separate trials, $P_{ET}CO_2$ was either permitted to fluctuate spontaneously (control trial) or was clamped at 1 mmHg above resting levels ($P_{ET}CO_2$ clamp trial). In the control trial, leg cycling with BFR decreased $P_{ET}CO_2$ ($\Delta -4.8 \pm 0.9$ mmHg vs. leg cycling exercise) secondary to hyperventilation, while ICA_Q , ICA_{CVC} , and MCA_{Vm} were unchanged, and MCA_{CVCi} decreased. However, in the $P_{ET}CO_2$ clamp trial, leg cycling with BFR increased both MCA_{Vm} ($\Delta 5.9 \pm 1.4$ cm \cdot s $^{-1}$) and ICA_Q ($\Delta 20.0 \pm 7.8$ ml \cdot min $^{-1}$), and attenuated the decrease in MCA_{CVCi} , while ICA_{CVC} was unchanged. In the control trial, PEI decreased $P_{ET}CO_2$ ($\Delta -7.0 \pm 1.3$ mmHg vs. rest), MCA_{Vm} and MCA_{CVCi} , whereas ICA_Q and ICA_{CVC} were unchanged. In contrast, in the $P_{ET}CO_2$ clamp trial both ICA_Q ($\Delta 18.5 \pm 11.9$ ml \cdot min $^{-1}$) and MCA_{Vm} ($\Delta 8.8 \pm 2.0$ cm \cdot s $^{-1}$) were elevated, while ICA_{CVC} and MCA_{CVCi} were unchanged. In conclusion, when hyperventilation-related decreases in $P_{ET}CO_2$ are prevented the activation of metabolically sensitive skeletal muscle afferent fibres increases cerebral blood flow.

New & Noteworthy: Muscle metaboreflex activation increases cerebral blood flow, but only when hyperventilation mediated reductions in the partial pressure of end-tidal carbon dioxide are prevented. These findings may have implications for individuals practicing exercise training with blood flow restriction and patient populations in whom exaggerated muscle metaboreflex sensitivity has been identified.

Key words: cerebrovascular circulation; exercise; metabolic activation

INTRODUCTION

Increases in cerebral blood flow during exercise are associated with upsurges in brain activation and metabolism within regions such as the motor-sensory cortex and supplementary motor area (23). Several other interacting mechanisms also contribute to the cerebral circulatory response accompanying exercise, including chemical, hemodynamic, autoregulatory and neural factors (42). The stimulation of group III and IV skeletal muscle afferents has also been implicated in the cerebral blood flow responses to exercise, but this remains incompletely understood (19, 20, 27). Group III and IV skeletal muscle afferents are responsive to metabolic (muscle metaboreflex) and mechanical (muscle mechanoreflex) perturbation. Alongside central command (feedforward signals from higher brain centres) and the arterial and cardiopulmonary baroreceptors, group III and IV skeletal muscle afferents play a key role in mediating the cardiovascular adjustments to exercise (18). Early studies established two approaches for the assessment of the muscle metaboreflex (3, 4, 52). The first approach involved blood flow restriction (BFR) to the exercising muscles using proximally placed inflatable occlusion cuffs, in order to create a mismatch between oxygen delivery and demand. This in turn evoked an accelerated accumulation of exercise-induced metabolites and enhanced activation of the metabolically sensitive skeletal muscle afferents. The second approach involved the complete circulatory arrest of the exercising skeletal muscle continuing into the recovery period while the muscle is quiescent (i.e., post-exercise ischemia; PEI). In this manner, metabolically sensitive skeletal muscle afferents may be activated in isolation by the trapping of exercise-induced metabolites within the muscle.

During the isolated activation of the muscle metaboreflex with PEI following handgrip, exercise-induced increases in middle cerebral artery (MCA) mean blood velocity (v_m) are not sustained and MCA_{v_m} returns to baseline (28, 46). This is in conflict with reports that the use of local anaesthesia to block sensory feedback from group III and IV skeletal muscle afferent fibres abolished the normal increase of MCA_{v_m} during static and dynamic handgrip (19, 20, 27). We recently observed that such contradictory reports may be attributable to muscle metaboreflex mediated increases in ventilation during PEI, which lead to a confounding reduction in the partial pressure of arterial carbon dioxide (P_aCO_2 ; indexed by the partial pressure of end-tidal carbon dioxide; $P_{ET}CO_2$) and a cerebral

vasoconstriction that prevents a muscle metaboreflex mediated increase in MCA_{Vm} . Indeed, the clamping of $P_{ET}CO_2$ at baseline values during PEI following fatiguing static handgrip resulted in an elevation in MCA_{Vm} (13).

The aforementioned studies possess limitations with regards to the assessment of cerebral blood flow and mode of muscle metaboreflex activation that we seek to address in the present investigation. First, for transcranial Doppler ultrasound measures of MCA_{Vm} to be representative of cerebral blood flow it must be assumed that MCA diameter remains constant. To circumvent this issue, direct measures of internal carotid artery (ICA) diameter (d), velocity (v_m) and thus ICA blood flow (ICA_Q) can be employed, but to date the contribution of skeletal muscle afferents to the ICA_Q responses to exercise remain unknown. Second, while the cerebral circulatory responses to muscle metaboreflex activation following exercise with PEI have been investigated, the responses to the enhanced muscle metaboreflex activation during exercise with BFR have not been considered. During PEI the muscle metaboreflex is activated in isolation from central command and skeletal muscle mechanoreflex (11, 16, 17). However, under clinical conditions such as peripheral vascular disease and chronic heart failure, where there is a hypoperfusion of the skeletal muscles, the muscle metaboreflex is not activated in isolation (7, 9, 24). Increasing metaboreflex signalling during exercise with BFR at the same time that central command and mechanoreflex are also activated would potentially provide a more realistic simulation of this paradigm. Furthermore, exercise with BFR is becoming an increasingly popular athletic training practice due to the potential for gains in muscle strength and endurance to occur without high-intensity training (54, 60). Spranger et al. (57) recently raised a ‘call for concern’ regarding the practice of exercise with BFR on the basis of the exaggerated increases in blood pressure and the associated risk of cardiovascular and cerebrovascular insult. At present the effects of BFR exercise on cerebral blood flow are unknown.

Given this background, we sought to determine; 1) the influence of the muscle metaboreflex activation on cerebral blood flow, 2) if changes in $P_{ET}CO_2$ are a key determinant of the cerebral blood flow response to muscle metaboreflex activation, and 3) whether the mode of muscle metaboreflex activation (i.e., during vs. following exercise) influences the corresponding cerebral blood flow response. To achieve this MCA_{Vm} and ICA_Q were measured during exercise with BFR to enhance

107 muscle metaboreflex activation and during isolated activation of the muscle metaboreflex with PEI.
108 Trials were conducted where $P_{ET}CO_2$ was permitted to fluctuate spontaneously and where $P_{ET}CO_2$ was
109 clamped at baseline values. We hypothesized that muscle metaboreflex activation would evoke an
110 increase in MCA_{Vm} and ICA_Q only when $P_{ET}CO_2$ was clamped at baseline.

METHODS

The study was approved by the Health, Safety and Ethics Committee of the School of Sport, Exercise and Rehabilitation Science at the University of Birmingham and was undertaken according to Declaration of Helsinki. Eleven male participants were recruited (age 25 ± 4 years; height 180 ± 1 cm; weight 71 ± 7 kg; mean \pm SD). After receiving a detailed verbal and written explanation of the experimental protocol, all participants signed the consent form. All participants were free of any cardiovascular, respiratory, neurological, renal or metabolic diseases and were not using any prescription or over-the-counter medication. Abstinence of caffeinate beverages, alcohol or exercise was requested 24 hours prior to experimental sessions. The room temperature was kept constant at 20-22°C, and external stimuli were kept to a minimum.

Measurements

Heart rate (HR) was monitored using lead II electrocardiogram and blood pressure was measured beat-to-beat from the middle finger of the right hand (Finometer Pro, Finapres Medical Systems, Arnhem, The Netherlands). Mean arterial pressure (MAP) was calculated offline by the integration of the arterial blood pressure waveform over a cardiac cycle. Resting blood pressure was verified by brachial artery blood pressure measurement made from the left arm using an automated sphygmomanometer (Tango+, SunTech Medical, USA). ICA_{Vm} and ICA_d were measured from the left side of the neck with duplex Doppler ultrasound (Logiq, GE Medical Systems, Milwaukee, USA) using a 10-MHz multifrequency linear-array transducer with a constant insonation angle of 60° relative to the skin. ICA measurements were performed 1 to 1.5 cm distal to the carotid bifurcation while the subject's chin was slightly elevated. To measure the ICA_d the brightness mode was used in a longitudinal section, the systolic and diastolic diameters were measured over 10 cardiac cycles, the mean diameter was calculated as: Mean diameter (cm) = [(systolic diameter \times 1/3)] + [(diastolic diameter \times 2/3)]. The Doppler velocity spectrum was analyzed using the pulsed wave mode and the time-averaged mean flow velocity obtained over 10 cardiac cycles. The internal carotid blood flow (ICA_Q ; ml·min⁻¹) was

calculated as $[ICA_{Vm} \times \pi \times (\text{diameter}/2)^2] \times 60$. ICA conductance (ICA_{CVC} ; $\text{ml} \cdot \text{min}^{-1} \cdot \text{mmHg}^{-1}$) was calculated as ICA_Q / MAP .

MCA_{Vm} was measured with a 2 MHz pulse wave transcranial Doppler ultrasound system (Doppler Box X; Compumedics Germany GmbH, Singen, Germany). The MCA was insonated via the temporal window above the zygomatic arch on the left side of the head. After finding a satisfactory signal, the probe was fixed in place with a headband and ultrasonic gel. The MCA vascular conductance index (MCA_{CVCi}) was calculated as MCA_{Vm} / MAP .

Participants wore a mouthpiece and nose-clip to permit breath-by-breath determination of minute ventilation (V_E) via a turbine volume transducer (VMM400; Interface Associates, Aliso Viejo, CA, USA). The end-tidal partial pressures of O_2 and CO_2 were determined using rapid response gas analyzers (Moxus Modular; AEI Technologies Inc, Pittsburg, USA). Analog data were digitally converted at 1 kHz and stored on a PC for offline analysis (Powerlab and LabChart Pro; ADInstruments, Dunedin, New Zealand).

Experimental Protocol

An initial familiarization session was first conducted where the participants experienced all the experimental methods and protocols. The experimental protocol was subsequently conducted over two laboratory visits separated by 3-7 days with the order of protocol 1 and 2 (day 1 or 2) decided according to a coin toss. For each protocol, two trials were performed. In one trial, $P_{ET}CO_2$ fluctuated normally while subjects breathed medical grade air, in another trial $P_{ET}CO_2$ was clamped at ~ 1 mmHg above the resting partial pressure. Trials were counterbalanced and separated by a minimum of 20 min. $P_{ET}CO_2$ clamping was undertaken using a dynamic end-tidal forcing system which uses a prediction – correction system, whereby $P_{ET}CO_2$ is controlled at the desired level by altering the composition of the inspired gas on a breath by breath basis (50).

Protocol I: Leg cycling with BFR

After instrumentation participants sat quietly in a semi-recumbent cycle ergometer and respiration was monitored for 10 min to determine the normal $P_{ET}CO_2$. Participants were then instructed to commence cycling exercise at 60 rpm. During the first ~3 to 5 min the workload was adjusted to reach a target HR of 120 beats·min⁻¹ after which ~10 min of steady-state cycling exercise was performed (Ex1). Following this period, bilateral thigh cuffs were inflated to 130 mmHg (Rapid Cuff Inflation System E20 AG101, Hokanson, Bellevue, USA) in order to partially restrict blood flow to exercising muscles and engage the muscle metaboreflex (BFR). The thigh cuffs were deflated after 3 min and a further 3 min of steady-state cycling exercise was performed under free-flow conditions (Ex2). ICA assessments were not carried out during Ex2. Ratings of perceived exertion (RPE) were obtained using the 1-10 Borg scale (12) at the end of the Ex1, BFR, and Ex2 periods. Mean HR, BP, respiratory and MCA_{Vm} data were obtained on a beat-to-beat basis and averages calculated at rest (3 min), Ex1 (last 1 min), BFR (last 1 min) and Ex2 (last 1 min). Ultrasound images for calculation of ICA_Q were obtained during the last 1 min of rest, last 1 min of Ex1 and last 1 min of leg cycling with BFR. Measurements were then pooled to provide a mean value for each experimental phase.

Protocol II: Leg cycling with PEI

As described above, following a 10 min rest period during which the normal $P_{ET}CO_2$ was determined, participants undertook cycling exercise on a semi-recumbent cycle ergometer (60 rpm). After a ~3-5 min period during which the workload was adjusted in order to reach the target HR of 120 beats·min⁻¹ participants undertook steady-state cycling exercise for 10 min (Ex). Fifteen seconds before the end of the exercise, bilateral thigh cuffs were inflated to 300 mmHg in order to occlude the blood flow to the exercising muscles and remained inflated for 3 min in order to isolate the activation of the muscle metaboreflex (PEI). RPE was obtained after 5 min of steady-state cycling exercise. Mean HR, BP, respiratory and MCA_{Vm} data were obtained on a beat-to-beat basis and averages calculated at rest (3 min), Ex (last 1 min) and PEI (last 1 min). Ultrasound images for calculation of

ICA_Q were obtained during last 1 min of rest, last 1 min of Ex and last 1 min of PEI. Measurements were then pooled to provide a mean value for each experimental phase.

Data and statistical analysis

Values are reported as means \pm SEM. Main effects of experimental phase (Rest, Ex, PEI or Rest, Ex1, BFR, Ex2), trial (control, P_{ET}CO₂ clamp) and interaction (phase x trial) were made using two-way repeated measures ANOVA followed by Student-Newman-Keuls post hoc test. Between trial comparisons of exercise workload were made using Student t-tests. Statistical significance was set to $p < 0.05$. Analyses were conducted using SigmaPlot 12.5 (Systat Software Inc, London, UK).

RESULTS

Leg cycling with BFR

The exercise workload was not different between trials (control 90 ± 9 W and $P_{ET}CO_2$ clamp trial 85 ± 9 W; $P > 0.05$). In the control trial, $P_{ET}CO_2$ was slightly increased from rest during leg cycling (Ex1 $\Delta 2.2 \pm 0.3$ mmHg, $P < 0.05$), decreased with BFR ($\Delta -4.8 \pm 0.9$ mmHg, $P < 0.05$), and returned to resting values upon the cessation of BFR (Ex2 $\Delta 0.8 \pm 0.5$ mmHg, $P > 0.05$ vs. Rest; Figure 1). By design, in the clamp trial $P_{ET}CO_2$ remained unchanged from the rest throughout all experimental phases. Leg cycling evoked similar increases in MCA_{V_m} during the control and $P_{ET}CO_2$ clamp trials ($P < 0.05$ Rest vs. Ex1; Figure 1). In the control trial, no change in MCA_{V_m} was observed during exercise with BFR, whereas in the $P_{ET}CO_2$ clamp trial MCA_{V_m} was increased ($P < 0.05$ vs. Rest, Ex1, and between conditions). In both trials, MCA_{V_m} was not different during leg cycling before and after BFR ($P > 0.05$ Ex1 vs. Ex2). MCA_{CV_i} was unchanged from rest during leg cycling ($P > 0.05$, Ex1 vs. Rest, Ex2 vs. Rest), but was decreased with BFR ($P < 0.05$ vs. Rest and Ex1; Table 1). The magnitude of this decrease was greater in the control trial than in the $P_{ET}CO_2$ clamp trial. ICA_Q was unchanged from rest to leg cycling (Ex1) in both trials, but was increased with BFR in the $P_{ET}CO_2$ clamp trial ($P < 0.05$ vs. Rest and between trials; Figure 1), secondary to an increase in ICA_{V_m} . ICA_d was unchanged throughout all experimental phases in both trials (Table 1). ICA_{CVC} was not different between trials and was similarly decreased from rest during leg cycling ($P < 0.05$ Rest vs. Ex1) then further decreased during leg cycling with BFR ($P < 0.05$ Ex1 vs. Rest) in both trials.

HR was not different between the control and $P_{ET}CO_2$ clamp trials at any experimental phase (Table 1). MAP was slightly, but higher in the $P_{ET}CO_2$ clamp trial (Figure 1). Leg cycling evoked increases in MAP, HR in both trials ($P < 0.05$ Ex1 vs. Rest), which were further increased during BFR ($P < 0.05$ vs. Ex1). During leg cycling following BFR, MAP and HR returned to values observed during leg cycling prior to BFR (Ex1 vs. Ex2, $P > 0.05$). RPE was not different between the control trial [Ex1 median, 4 (interquartile range, 4-5), BFR, 8 (7-8) and Ex2 5 (4-6); $P < 0.05$] and the $P_{ET}CO_2$ clamp trial [Ex1 4 (3-6), BFR 8, (7-8), Ex2 5 (4-7); $P < 0.05$] (Wilcoxon signed-rank test).

Leg cycling with PEI

Exercise workload was not different between trials (control 89 ± 9 W and $P_{ET}CO_2$ clamp trial 89 ± 9 W; $P > 0.05$). In the control trial, $P_{ET}CO_2$ was unchanged from rest during leg cycling and decreased with PEI ($\Delta -7 \pm 1$ mmHg, $P < 0.05$; Figure 2), but by design, in the $P_{ET}CO_2$ clamp trial it remained not different from rest throughout all experimental phases. In both the control and $P_{ET}CO_2$ clamp trials, MCA_{Vm} increased from rest during leg cycling ($P \leq 0.05$; Figure 2). During PEI, in the control trial MCA_{Vm} was not different from rest ($P > 0.05$), whereas in the $P_{ET}CO_2$ clamp trial MCA_{Vm} remained elevated ($P < 0.05$ vs. rest and between trials). In the control trial, MCA_{CVCi} was not different from rest during leg cycling and was decreased from rest during PEI ($P < 0.05$ vs. Rest and between trials; Table 2). MCA_{CVCi} was not different from rest throughout $P_{ET}CO_2$ clamp trial ($P > 0.05$; Table 2). ICA_Q was not different from rest during leg cycling in both control and $P_{ET}CO_2$ clamp trials ($P > 0.05$; Figure 2). However, during PEI, ICA_Q was higher in the $P_{ET}CO_2$ clamp trial than in the control trial ($P < 0.05$; Figure 2). ICA_d was not different throughout all experimental phases in both trials (Table 2). ICA_{CVC} was greater in the $P_{ET}CO_2$ clamp trial, but decreased similarly from rest during leg cycling ($P < 0.05$ vs. Rest) in both trials. During PEI, ICA_{CVC} remained at levels sustained during exercise ($P < 0.05$ vs. Rest, $P > 0.05$ vs. Ex) in both trials. MAP and HR (Table 2; Figure 2) were not different between the control and $P_{ET}CO_2$ clamp trials at any experimental phase. Leg cycling evoked increases in MAP and HR in both trials ($P < 0.05$ vs. rest). In both trials, MAP was further increased with PEI from the level observed during leg cycling ($P < 0.05$ vs. Rest and Ex), whereas HR fell, but remained above resting levels ($P < 0.05$ vs. Rest and Ex). RPE were not different during leg cycling in the control [median, 5 (interquartile range, 4-7)] and the $P_{ET}CO_2$ clamp trials [5 (4-6)] (Wilcoxon signed-rank test).

DISCUSSION

The major novel finding of this study is that the muscle metaboreflex failed to elevate either MCA_{V_m} or ICA_Q , when engaged by leg cycling with BFR or isolated during PEI following leg cycling under control conditions. However, a significant reduction in $P_{ET}CO_2$, secondary to an increase in V_E , was induced by muscle metaboreflex activation with either BFR or PEI. Accordingly, when $P_{ET}CO_2$ was clamped at resting levels muscle metaboreflex-mediated increases in MCA_{V_m} and ICA_Q were revealed during both BFR and PEI. Thus, in accordance with our original hypothesis, these findings demonstrate that when hyperventilation-related decreases in $P_{ET}CO_2$ are prevented the muscle metaboreflex increases cerebral blood flow, and this occurs irrespective of the mode of muscle metaboreflex activation.

A potential explanation why exercise with BFR, or indeed PEI, does not increase cerebral perfusion may be that the activation of metabolically sensitive skeletal muscle afferents evokes an increase in ventilation (5), which leads to a confounding reduction in P_aCO_2 (indexed by $P_{ET}CO_2$). The contribution of group III and IV skeletal muscle afferents to the control of breathing remains controversial, nevertheless in agreement with several previous reports we observed an increase in V_E during muscle metaboreflex activation (1, 13, 15, 44) and a reduction in $P_{ET}CO_2$. CO_2 is a powerful dilator of the cerebral vasculature (31) and decreases in P_aCO_2 lead to cerebral vasoconstriction (2). In the present study, the clamping of $P_{ET}CO_2$ at resting levels unmasked a muscle metaboreflex-mediated increase in MCA_{V_m} and ICA_Q during leg cycling with BFR and during PEI following leg cycling. Furthermore, the degree of cerebral vasoconstriction (i.e., the magnitude of the reduction in MCA_{CVci}) during PEI and exercise with BFR was attenuated in the $P_{ET}CO_2$ clamp trial, although a similar effect was not observed for ICA_{CVC} . These observations are in concordance with our earlier report that PEI following fatiguing ischemic handgrip exercise only increases MCA_{V_m} when $P_{ET}CO_2$ is clamped at resting levels (13). Such observations may help to explain why others have not shown MCA_{V_m} to be elevated during PEI. Indeed, Jorgensen et al., (28) also reported that P_aCO_2 was decreased below resting levels during PEI following cycling exercise. In agreement with Friedman et al., (19, 20) and Jorgensen et al., (29), who demonstrated that pharmacological blockade of sensory feedback from skeletal muscle afferents diminished the increase in cerebral perfusion during exercise, the results of the present study

support a role for the muscle metaboreflex in the regulation of cerebral blood flow during exercise. This may be attributable to the pairing of local neuronal activation and perfusion (i.e., neural-vascular coupling), that is to say cerebral flow increases in order to increase O₂ delivery in accordance with increased metabolic. Studies employing advanced imaging techniques have shown that isolated metaboreflex stimulation (PEI) evokes increased activity in discrete brain regions (e.g., medial and lateral dorsal medulla, contralateral insula, primary and secondary somatosensory cortex) (53). It is acknowledged that part of the cerebrovascular responses to PEI arises on account of the discomfort associated with this manoeuvre (35). In the present study we are not able to quantify the contribution of local discomfort to the cerebrovascular responses to PEI and BFR, but note that without the clamping of P_{ET}CO₂ at resting levels no changes in cerebral perfusion were observed. As such, the combination of P_{ET}CO₂ clamping and brain imaging modalities may provide additional insights into the effects of muscle metaboreflex activation on regional brain activation. The influence of exercise-induced increases of MAP on cerebral blood flow is somewhat controversial. The observation that during PEI, MCA_{Vm} remains at resting levels while MAP is elevated is part of the reason why the direct influence of blood pressure on the exercise-induced increase of cerebral perfusion has on occasion been discounted (26, 47, 55). However, when considering the effects of MAP on cerebral perfusion during exercise the potentially confounding effects of changes in P_{ET}CO₂ should be considered. With P_aCO₂ controlled, MCA_{Vm} changes by ~0.8% per mmHg change in MAP within the so called “autoregulatory range” (34). We observed that, BFR evoked a 44 mmHg increase from rest in MAP and a 26% increase in MCA_{Vm}, while during PEI, MAP was elevated by 29 mmHg and MCA_{Vm} elevated by 17%. As such, increases in MAP may reasonably be expected to contribute, at least in part, for the observed increase in cerebral blood flow. The ventilatory response to leg cycling with BFR was enhanced when P_{ET}CO₂ was clamped. As neither the exercise workload nor the thigh cuff pressure were different between conditions, it is unlikely that this is attributable to a difference in the degree of muscle afferent activation within the active skeletal muscles. However, in the absence of a direct assessment of local metabolites this possibility cannot be excluded. Alternatively, the clamping of P_{ET}CO₂ may have eliminated part of the chemoreflex-mediated inhibition of the ventilatory response. It is possible that the greater ventilatory

response to exercise observed when $P_{ET}CO_2$ was clamped, could evoke a respiratory muscle metaboreflex and augment the concomitant blood pressure response (51). However, an exaggerated increase in blood pressure did not accompany the greater ventilatory response to exercise with BFR in the clamp trial.

The present study extends earlier reports examining the contribution of skeletal muscle afferents to cerebral blood flow regulation in two important ways. Rather than relying on transcranial Doppler ultrasound measures of MCA_{Vm} to estimate cerebral perfusion, we have used duplex Doppler ultrasound measures of cerebral blood flow (ICA_Q). In order that transcranial Doppler ultrasound measures of MCA_{Vm} are representative of cerebral blood flow, it must be assumed that MCA_d remains constant. As such, this is the first study to determine the influence of the muscle metaboreflex activation on cerebral blood flow in humans. In addition to use of PEI to assess the effect of isolated skeletal muscle metaboreflex activation on cerebral blood flow, exercise with BFR was undertaken. As indicated above, this has applied relevance to those undertaking such practices to induce athletic enhancements, but also patient populations in whom muscle metaboreflex activation may be heightened during exercise as a consequence of skeletal muscle under-perfusion (e.g., chronic heart failure, peripheral vascular disease). Despite exercise with BFR and PEI evoking similar reductions from rest in $P_{ET}CO_2$ (Δ -2.6 and Δ -5.0 mmHg for BFR and PEI, respectively), MCA_{Vm} was significantly elevated from rest during exercise with BFR such that it was similar to that observed during free-flow exercise, whereas MCA_{Vm} was not different from rest during PEI. Such findings may be explained by a greater elevations in cardiac output and the concomitant activation of central command during exercise with BFR compared to PEI (36). Nevertheless, independent of the mode of muscle metaboreflex activation, when muscle metaboreflex mediated reductions in $P_{ET}CO_2$ were prevented increases in cerebral blood flow were observed.

Regular exercise with BFR (e.g., Kaatsu training) has been reported to enhance cardiorespiratory fitness (60) and skeletal muscle mass in healthy young and older individuals (32, 33, 63, 64), and patients (56). It is believed that exercise training with BFR exaggerates the normal accumulation of metabolites within the active skeletal muscle, thus promoting muscle growth and force generating capacity without the need for high-intensity training (21, 22, 43, 62). In a recent article, Spranger et al.,

(57) raised a ‘call for concern’ about this practice on the basis that it engages the muscle metaboreflex, which is known to powerfully increase sympathetic nerve activity to the heart and peripheral vasculature and can inhibit cardiac parasympathetic activity (18). As a consequence of these autonomic alterations, exercise with BFR evokes pronounced increases in peripheral vascular resistance, cardiac output and blood pressure (59). This could be of particular concern to patients in whom exaggerated skeletal muscle afferent sensitivity has been identified (e.g., hypertension, chronic heart failure, chronic obstructive pulmonary disease, type 2 diabetes) (7, 9, 24, 25, 45) and could raise the risk of cardiovascular and cerebrovascular events (57). Indeed, in the present study of healthy individuals, exercise with BFR raised MAP by ~ 27 mmHg and heart rate by ~ 22 beats \cdot min $^{-1}$ from levels established during a preceding period of leg cycling under free-flow conditions. However, despite such hemodynamic alterations cerebral perfusion was unchanged during exercise with BFR, thus calling into question the contention that exercise with BFR may increase the risk of cerebrovascular injury as a consequence of a large local hyperaemic response. In fact we observed that exercise with BFR evoked a lower than expected cerebral blood flow, on account of the coincident hyperventilation and hypocapnia linked cerebral vasoconstriction. During high intensity dynamic exercise, particularly when combined with hypoxia, a reduced cerebral oxygenation has been postulated as a fatigue mechanism (8, 40, 55). As such, in patients who exhibit exaggerated skeletal muscle afferent feedback and an excessive hyperventilatory response to exercise (e.g., chronic heart failure, congestive obstructive pulmonary disease) (7, 9, 58), the associated reduction in $P_a\text{CO}_2$ and cerebral vasoconstriction may precipitate exercise intolerance via a central mechanism (41, 48). Future studies utilizing arterial-internal jugular venous blood sampling are required, particularly in chronic disease populations, to better understand how cerebral metabolism (e.g., O_2 delivery, cerebral metabolic rate for O_2 , fractional O_2 extraction) is affected by the activation of skeletal muscle afferents while $P_{\text{ET}}\text{CO}_2$ is clamped at baseline levels.

There are several limitations on the present study, $P_{\text{ET}}\text{CO}_2$ was used as a surrogate for $P_a\text{CO}_2$, although during exercise $P_{\text{ET}}\text{CO}_2$ could have overestimated $P_a\text{CO}_2$ (49) there is a strong correlation between $P_{\text{ET}}\text{CO}_2$ and $P_a\text{CO}_2$ across all levels of physiologic dead space (37). We used external thigh cuffs to decrease/occlude blood flow to the legs, although no direct measures of the leg blood flow or

local metabolite concentrations were possible, we observed marked increases in MAP during this manoeuvre indicative of muscle metaboreflex activation, which strongly suggest that reductions in blood flow were successfully induced. BFR may have increased in central command, which could have contributed to the increase of cerebral blood flow (6). Indeed, increases in RPE, historically related to central command (38, 39), were noted during leg cycling with BFR. Human studies are needed in which the cardiovascular and cerebrovascular responses to the application of BFR during leg cycling are evaluated before and after pharmacological inhibition of feedback from group III and IV skeletal muscle afferents (e.g., intrathecal fentanyl) in order to test the contribution of central command to this manoeuvre. Automated edge-tracking software was not used in the present study and this may be a limitation (61), although intraclass correlations between repeated rest and exercise measures made during a study visit were high (i.e., >0.8). Finally, we have only evaluated the effect of skeletal muscle afferent feedback with BFR during moderate cycling exercise, and care should be taken when directly extrapolating our findings to other exercise intensities and modalities. This is important in light of the concept that athletes may utilise BFR training in order to try to obtain a desirable training effects but at a lower exercise intensity (21, 22, 43, 62).

In conclusion, the findings of the present study indicate that only when hyperventilation-related decreases in $P_{ET}CO_2$ are prevented does the activation of metabolically sensitive skeletal muscle afferent fibres evoke an increase in cerebral blood flow, irrespective of the mode of activation (i.e. during or following Ex).

Acknowledgements

The authors appreciate the time and effort expended by the participants in this study, and would like to thank Joel Zvick and Prokopios Barmparigos for their technical support.

Sources of funding

381 This study was supported by Coordination for the Improvement of Higher Education Personnel
382 (CAPES, Brazil) through the program Science without Borders.

383

384 Conflict of interest

385 None to declare.

1. **Adams L, Guz A, Innes JA, and Murphy K.** The early circulatory and ventilatory response to voluntary and electrically induced exercise in man. *The Journal of physiology* 383: 19-30, 1987.
2. **Ainslie PN, and Duffin J.** Integration of cerebrovascular CO₂ reactivity and chemoreflex control of breathing: mechanisms of regulation, measurement, and interpretation. *American journal of physiology Regulatory, integrative and comparative physiology* 296: R1473-1495, 2009.
3. **Alam M, and Smirk FH.** Observations in man on a pulse-accelerating reflex from the voluntary muscles of the legs. *The Journal of physiology* 92: 167-177, 1938.
4. **Alam M, and Smirk FH.** Observations in man upon a blood pressure raising reflex arising from the voluntary muscles. *The Journal of physiology* 89: 372-383, 1937.
5. **Amann M, Blain GM, Proctor LT, Sebranek JJ, Pegelow DF, and Dempsey JA.** Group III and IV muscle afferents contribute to ventilatory and cardiovascular response to rhythmic exercise in humans. *Journal of applied physiology* 109: 966-976, 2010.
6. **Amann M, Proctor LT, Sebranek JJ, Pegelow DF, and Dempsey JA.** Opioid-mediated muscle afferents inhibit central motor drive and limit peripheral muscle fatigue development in humans. *The Journal of physiology* 587: 271-283, 2009.
7. **Amann M, Regan MS, Kobitarty M, Eldridge MW, Boutellier U, Pegelow DF, and Dempsey JA.** Impact of pulmonary system limitations on locomotor muscle fatigue in patients with COPD. *American journal of physiology Regulatory, integrative and comparative physiology* 299: R314-324, 2010.
8. **Amann M, Sidhu SK, Weavil JC, Mangum TS, and Venturelli M.** Autonomic responses to exercise: group III/IV muscle afferents and fatigue. *Autonomic neuroscience : basic & clinical* 188: 19-23, 2015.
9. **Amann M, Venturelli M, Ives SJ, Morgan DE, Gmelch B, Witman MA, Jonathan Groot H, Walter Wray D, Stehlik J, and Richardson RS.** Group III/IV muscle afferents impair limb blood in patients with chronic heart failure. *International journal of cardiology* 174: 368-375, 2014.
10. **Bolduc V, Thorin-Trescases N, and Thorin E.** Endothelium-dependent control of cerebrovascular functions through age: exercise for healthy cerebrovascular aging. *American journal of physiology Heart and circulatory physiology* 305: H620-633, 2013.
11. **Bonde-Petersen F, Rowell LB, Murray RG, Blomqvist GG, White R, Karlsson E, Campbell W, and Mitchell JH.** Role of cardiac output in the pressor responses to graded muscle ischemia in man. *Journal of applied physiology: respiratory, environmental and exercise physiology* 45: 574-580, 1978.
12. **Borg GA.** Psychophysical bases of perceived exertion. *Medicine and science in sports and exercise* 14: 377-381, 1982.
13. **Braz ID, Scott C, Simpson LL, Springham EL, Tan BW, Balanos GM, and Fisher JP.** Influence of muscle metaboreceptor stimulation on middle cerebral artery blood velocity in humans. *Experimental physiology* 99: 1478-1487, 2014.
14. **Davenport MH, Hogan DB, Eskes GA, Longman RS, and Poulin MJ.** Cerebrovascular reserve: the link between fitness and cognitive function? *Exercise and sport sciences reviews* 40: 153-158, 2012.
15. **Duncan G, Johnson RH, and Lambie DG.** Role of sensory nerves in the cardiovascular and respiratory changes with isometric forearm exercise in man. *Clinical science* 60: 145-155, 1981.
16. **Fisher JP, Adlan AM, Shantsila A, Secher JF, Sorensen H, and Secher NH.** Muscle metaboreflex and autonomic regulation of heart rate in humans. *The Journal of physiology* 591: 3777-3788, 2013.
17. **Fisher JP, Seifert T, Hartwich D, Young CN, Secher NH, and Fadel PJ.** Autonomic control of heart rate by metabolically sensitive skeletal muscle afferents in humans. *The Journal of physiology* 588: 1117-1127, 2010.
18. **Fisher JP, Young CN, and Fadel PJ.** Autonomic adjustments to exercise in humans. *Comprehensive Physiology* 5: 475-512, 2015.
19. **Friedman DB, Friberg L, Mitchell JH, and Secher NH.** Effect of axillary blockade on regional cerebral blood flow during static handgrip. *Journal of applied physiology* 71: 651-656, 1991.

20. **Friedman DB, Friberg L, Payne G, Mitchell JH, and Secher NH.** Effects of axillary blockade on regional cerebral blood flow during dynamic hand contractions. *Journal of applied physiology* 73: 2120-2125, 1992.
21. **Fry CS, Glynn EL, Drummond MJ, Timmerman KL, Fujita S, Abe T, Dhanani S, Volpi E, and Rasmussen BB.** Blood flow restriction exercise stimulates mTORC1 signaling and muscle protein synthesis in older men. *Journal of applied physiology* 108: 1199-1209, 2010.
22. **Fujita S, Abe T, Drummond MJ, Cadenas JG, Dreyer HC, Sato Y, Volpi E, and Rasmussen BB.** Blood flow restriction during low-intensity resistance exercise increases S6K1 phosphorylation and muscle protein synthesis. *Journal of applied physiology* 103: 903-910, 2007.
23. **Gross PM, Marcus ML, and Heistad DD.** Regional distribution of cerebral blood flow during exercise in dogs. *Journal of applied physiology: respiratory, environmental and exercise physiology* 48: 213-217, 1980.
24. **Haas TL, Lloyd PG, Yang HT, and Terjung RL.** Exercise training and peripheral arterial disease. *Comprehensive Physiology* 2: 2933-3017, 2012.
25. **Holwerda SW, Restaino RM, Acevedo CM, Lastra G, Fisher JP, and Fadel PJ.** Augmented pressor and sympathetic responses to skeletal muscle metaboreflex activation in type 2 diabetes patients. *American journal of physiology Heart and circulatory physiology* 2015.
26. **Ide K, and Secher NH.** Cerebral blood flow and metabolism during exercise. *Progress in neurobiology* 61: 397-414, 2000.
27. **Jorgensen LG, Perko G, Payne G, and Secher NH.** Effect of limb anesthesia on middle cerebral response to handgrip. *The American journal of physiology* 264: H553-559, 1993.
28. **Jorgensen LG, Perko G, and Secher NH.** Regional cerebral artery mean flow velocity and blood flow during dynamic exercise in humans. *Journal of applied physiology* 73: 1825-1830, 1992.
29. **Jorgensen LG, Perko M, Perko G, and Secher NH.** Middle cerebral artery velocity during head-up tilt induced hypovolaemic shock in humans. *Clinical physiology* 13: 323-336, 1993.
30. **Joyner MJ.** Buying into healthy blood vessels: exercise and aging. *Journal of applied physiology* 117: 421-422, 2014.
31. **Kety SS, and Schmidt CF.** The Effects of Altered Arterial Tensions of Carbon Dioxide and Oxygen on Cerebral Blood Flow and Cerebral Oxygen Consumption of Normal Young Men. *The Journal of clinical investigation* 27: 484-492, 1948.
32. **Lixandrao ME, Ugrinowitsch C, Laurentino G, Libardi CA, Aihara AY, Cardoso FN, Tricoli V, and Roschel H.** Effects of exercise intensity and occlusion pressure after 12 weeks of resistance training with blood-flow restriction. *European journal of applied physiology* 2015.
33. **Loenneke JP, Wilson JM, Marin PJ, Zourdos MC, and Bemben MG.** Low intensity blood flow restriction training: a meta-analysis. *European journal of applied physiology* 112: 1849-1859, 2012.
34. **Lucas SJ, Tzeng YC, Galvin SD, Thomas KN, Ogoh S, and Ainslie PN.** Influence of changes in blood pressure on cerebral perfusion and oxygenation. *Hypertension* 55: 698-705, 2010.
35. **Macefield VG, and Henderson LA.** Autonomic responses to exercise: cortical and subcortical responses during post-exercise ischaemia and muscle pain. *Autonomic neuroscience : basic & clinical* 188: 10-18, 2015.
36. **McNulty CL, Moody WE, Wagenmakers AJ, and Fisher JP.** Effect of muscle metaboreflex activation on central hemodynamics and cardiac function in humans. *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme* 39: 861-870, 2014.
37. **McSwain SD, Hamel DS, Smith PB, Gentile MA, Srinivasan S, Meliones JN, and Cheifetz IM.** End-tidal and arterial carbon dioxide measurements correlate across all levels of physiologic dead space. *Respiratory care* 55: 288-293, 2010.
38. **Mitchell JH.** Cardiovascular control during exercise: central and reflex neural mechanisms. *The American journal of cardiology* 55: 34D-41D, 1985.
39. **Mitchell JH.** Neural circulatory control during exercise: early insights. *Experimental physiology* 98: 867-878, 2013.
40. **Nielsen HB, Boushel R, Madsen P, and Secher NH.** Cerebral desaturation during exercise reversed by O2 supplementation. *The American journal of physiology* 277: H1045-1052, 1999.

41. **Nybo L, and Rasmussen P.** Inadequate cerebral oxygen delivery and central fatigue during strenuous exercise. *Exercise and sport sciences reviews* 35: 110-118, 2007.
42. **Ogoh S, and Ainslie PN.** Cerebral blood flow during exercise: mechanisms of regulation. *Journal of applied physiology* 107: 1370-1380, 2009.
43. **Pearson SJ, and Hussain SR.** A review on the mechanisms of blood-flow restriction resistance training-induced muscle hypertrophy. *Sports medicine* 45: 187-200, 2015.
44. **Piepoli M, Ponikowski P, Clark AL, Banasiak W, Capucci A, and Coats AJ.** A neural link to explain the "muscle hypothesis" of exercise intolerance in chronic heart failure. *American heart journal* 137: 1050-1056, 1999.
45. **Poole DC, Hirai DM, Copp SW, and Musch TI.** Muscle oxygen transport and utilization in heart failure: implications for exercise (in)tolerance. *American journal of physiology Heart and circulatory physiology* 302: H1050-1063, 2012.
46. **Pott F, Ray CA, Olesen HL, Ide K, and Secher NH.** Middle cerebral artery blood velocity, arterial diameter and muscle sympathetic nerve activity during post-exercise muscle ischaemia. *Acta physiologica Scandinavica* 160: 43-47, 1997.
47. **Querido JS, and Sheel AW.** Regulation of cerebral blood flow during exercise. *Sports medicine* 37: 765-782, 2007.
48. **Rasmussen P, Nielsen J, Overgaard M, Krogh-Madsen R, Gjedde A, Secher NH, and Petersen NC.** Reduced muscle activation during exercise related to brain oxygenation and metabolism in humans. *The Journal of physiology* 588: 1985-1995, 2010.
49. **Robbins PA, Conway J, Cunningham DA, Khamnei S, and Paterson DJ.** A comparison of indirect methods for continuous estimation of arterial PCO₂ in men. *Journal of applied physiology* 68: 1727-1731, 1990.
50. **Robbins PA, Swanson GD, Micco AJ, and Schubert WP.** A fast gas-mixing system for breath-to-breath respiratory control studies. *Journal of applied physiology: respiratory, environmental and exercise physiology* 52: 1358-1362, 1982.
51. **Romer LM, Lovering AT, Haverkamp HC, Pegelow DF, and Dempsey JA.** Effect of inspiratory muscle work on peripheral fatigue of locomotor muscles in healthy humans. *The Journal of physiology* 571: 425-439, 2006.
52. **Rowell LB, Sheriff DD, Wyss CR, and Scher AM.** The nature of the exercise stimulus. *Acta physiologica Scandinavica Supplementum* 556: 7-14, 1986.
53. **Sander M, Macefield VG, and Henderson LA.** Cortical and brain stem changes in neural activity during static handgrip and postexercise ischemia in humans. *Journal of applied physiology* 108: 1691-1700, 2010.
54. **Scott BR, Loenneke JP, Slaterry KM, and Dascombe BJ.** Exercise with blood flow restriction: an updated evidence-based approach for enhanced muscular development. *Sports medicine* 45: 313-325, 2015.
55. **Secher NH, Seifert T, and Van Lieshout JJ.** Cerebral blood flow and metabolism during exercise: implications for fatigue. *Journal of applied physiology* 104: 306-314, 2008.
56. **Segal NA, Williams GN, Davis MC, Wallace RB, and Mikesky AE.** Efficacy of blood flow-restricted, low-load resistance training in women with risk factors for symptomatic knee osteoarthritis. *PM R* 7: 376-384, 2015.
57. **Spranger MD, Krishnan AC, Levy PD, O'Leary DS, and Smith SA.** Blood flow restriction training and the exercise pressor reflex: a call for concern. *American journal of physiology Heart and circulatory physiology* ajpheart 00208 02015, 2015.
58. **Stone AJ, and Kaufman MP.** The exercise pressor reflex and peripheral artery disease. *Autonomic neuroscience : basic & clinical* 188: 69-73, 2015.
59. **Sugawara J, Tomoto T, and Tanaka H.** Impact of leg blood flow restriction during walking on central arterial hemodynamics. *American journal of physiology Regulatory, integrative and comparative physiology* 309: R732-739, 2015.
60. **Taylor CW, Ingham SA, and Ferguson RA.** Acute and chronic effect of sprint interval training combined with post-exercise blood flow restriction in trained individuals. *Experimental physiology* 2015.

- 544 61. **Thomas KN, Lewis NC, Hill BG, and Ainslie PN.** Technical recommendations for the use of
545 carotid duplex ultrasound for the assessment of extracranial blood flow. *American journal of physiology*
546 *Regulatory, integrative and comparative physiology* 309: R707-720, 2015.
- 547 62. **Wernbom M, Apro W, Paulsen G, Nilsen TS, Blomstrand E, and Raastad T.** Acute low-load
548 resistance exercise with and without blood flow restriction increased protein signalling and number of
549 satellite cells in human skeletal muscle. *European journal of applied physiology* 113: 2953-2965, 2013.
- 550 63. **Yasuda T, Fukumura K, Fukuda T, Iida H, Imuta H, Sato Y, Yamasoba T, and Nakajima**
551 **T.** Effects of low-intensity, elastic band resistance exercise combined with blood flow restriction on
552 muscle activation. *Scandinavian journal of medicine & science in sports* 24: 55-61, 2014.
- 553 64. **Yasuda T, Fukumura K, Uchida Y, Koshi H, Iida H, Masamune K, Yamasoba T, Sato Y,**
554 **and Nakajima T.** Effects of Low-Load, Elastic Band Resistance Training Combined With Blood Flow
555 Restriction on Muscle Size and Arterial Stiffness in Older Adults. *The journals of gerontology Series A,*
556 *Biological sciences and medical sciences* 70: 950-958, 2015.

557

FIGURE LEGENDS

Figure 1. Cardiorespiratory and cerebrovascular responses to leg cycling under free-flow conditions (Ex1, Ex2) and with blood flow restriction (BFR). Partial pressure of end-tidal carbon dioxide ($P_{ET}CO_2$); middle cerebral artery blood velocity (MCA_{Vm}); internal carotid artery blood flow (ICA_Q); mean arterial pressure (MAP). Ex, exercise; BFR, blood flow restriction. Values are mean \pm SEM. * $P<0.05$ vs. rest, $\dagger P<0.05$ vs. Ex1, $\ddagger P<0.05$ vs. control.

Figure 2. Cardiorespiratory and cerebrovascular responses to leg cycling and post-exercise ischemia (PEI). Partial pressure of end-tidal carbon dioxide ($P_{ET}CO_2$); middle cerebral artery blood velocity (MCA_{Vm}); internal carotid artery blood flow (ICA_Q); mean arterial pressure (MAP). Ex, exercise; PEI, post exercise ischemia. Values are mean \pm SEM. * $P<0.05$ vs. rest, $\dagger P<0.05$ vs. Ex, $\ddagger P<0.05$ vs. control.

Control	3.3 ± 0.2	2.9 ± 0.2	2.0 ± 0.1				
P _{ET} CO ₂ Clamp	3.2 ± 0.2	2.8 ± 0.2	2.4 ± 0.1		<0.001	0.321	0.108
SBP (mmHg)							
Control	117 ± 2	153 ± 7	191 ± 8	146 ± 7			
P _{ET} CO ₂ Clamp	115 ± 3	153 ± 5	195 ± 6	146 ± 6	<0.001	0.611	0.402
DBP (mmHg)							
Control	67 ± 3	72 ± 3	96 ± 3	75 ± 4			
P _{ET} CO ₂ Clamp	70 ± 2	79 ± 3	101 ± 5	80 ± 3	<0.001	<0.001	0.522
HR (beats·min ⁻¹)							
Control	64 ± 3	122 ± 1	144 ± 3	124 ± 2			
P _{ET} CO ₂ Clamp	65 ± 3	121 ± 1	145 ± 2	125 ± 2	<0.001	0.450	0.760

Values are mean±SEM. V_E, ventilation; MCA_{CVCI}, middle cerebral artery conductance; ICA_{Vm}, internal carotid artery mean velocity; ICA_d, internal carotid artery diameter; ICA_{CVCI}, internal carotid artery conductance; SBP, systolic blood pressure; MAP, mean arterial pressure; HR, heart rate; Ex, exercise; BFR, blood flow restriction. Values are mean ± SEM. * P<0.05 vs. Rest, †P<0.05 vs. Ex1, ‡P<0.05 vs. Control.

Table 2. Cardiorespiratory and cerebrovascular responses to leg cycling and post-exercise ischemia (PEI)

	Experimental Phase			P value		
	Rest	Ex	PEI	Phase	Condition	Interaction
V _E (l·min ⁻¹)						
Control	9 ± 1	40 ± 2*	16 ± 3†	<0.001	0.011	0.001
P _{ET} CO ₂ Clamp	13 ± 1	40 ± 3*	25 ± 3*†‡			
MCA _{CVCi} (cm·s ⁻¹ ·mmHg ⁻¹)						
Control	0.60 ± 0.05	0.55 ± 0.03	0.43 ± 0.03*†	0.001	0.004	0.031
P _{ET} CO ₂ Clamp	0.64 ± 0.04	0.59 ± 0.04	0.56 ± 0.04‡			
ICA _{Vm} (cm·s ⁻¹)						
Control	25 ± 1	26 ± 1	23 ± 2	0.735	<0.001	0.019
P _{ET} CO ₂ Clamp	27 ± 1	28 ± 1	29 ± 2‡			
ICA _d (cm)						
Control	0.45 ± 0.01	0.45 ± 0.01	0.45 ± 0.01	0.306	0.572	0.784
P _{ET} CO ₂ Clamp	0.46 ± 0.01	0.45 ± 0.01	0.45 ± 0.01			
ICA _{CVC} (ml·min ⁻¹ ·mmHg ⁻¹)						
Control	2.8 ± 0.2	2.4 ± 0.2	1.9 ± 0.2	<0.001	<0.001	0.122

P _{ET} CO ₂ Clamp	3.1 ± 0.2	2.6 ± 0.1	2.5 ± 0.2			
SBP (mmHg)						
Control	118 ± 4	155 ± 7	145 ± 7			
P _{ET} CO ₂ Clamp	116 ± 4	159 ± 9	148 ± 8	<0.001	0.729	0.469
DBP (mmHg)						
Control	72 ± 3	83 ± 4	98 ± 5			
P _{ET} CO ₂ Clamp	69 ± 2	80 ± 3	94 ± 3	<0.001	0.157	0.932
HR (beats·min ⁻¹)						
Control	61 ± 3	121 ± 2	95 ± 5			
P _{ET} CO ₂ Clamp	63 ± 3	123 ± 1	93 ± 3	<0.001	0.764	0.416

Values are mean±SEM. V_E, ventilation; MCA_{CVC}, middle cerebral artery conductance; ICA_{Vm}, internal carotid artery mean velocity; ICA_d, internal carotid artery diameter; ICA_{CVCi}, internal carotid artery conductance; SBP, systolic blood pressure; MAP, mean arterial pressure; HR, heart rate; Ex, exercise; PEI, post exercise ischemia. Values are mean ± SEM. * P<0.05 vs. Rest, †P<0.05 vs. Ex, ‡P<0.05 vs. Control.

582

583

584

585

586

587

588

589

590

591



